


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



Volume 5. Issue 12. Pages 1847-2008. 2010
ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF**DR. PAWAN K AGRAWAL**

*Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us*

EDITORS**PROFESSOR ALESSANDRA BRACA**

*Dipartimento di Chimica Bioorganica e Biofarmacia,
Universita di Pisa,
via Bonanno 33, 56126 Pisa, Italy
braca@farm.unipi.it*

PROFESSOR DEAN GUO

*State Key Laboratory of Natural and Biomimetic Drugs,
School of Pharmaceutical Sciences,
Peking University,
Beijing 100083, China
gda5958@163.com*

PROFESSOR J. ALBERTO MARCO

*Departamento de Química Organica,
Universidade de Valencia,
E-46100 Burjassot, Valencia, Spain
alberto.marco@uv.es*

PROFESSOR YOSHIHIRO MIMAKI

*School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimakiy@ps.toyaku.ac.jp*

PROFESSOR STEPHEN G. PYNE

*Department of Chemistry
University of Wollongong
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au*

PROFESSOR MANFRED G. REINECKE

*Department of Chemistry,
Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu*

PROFESSOR WILLIAM N. SETZER

*Department of Chemistry
The University of Alabama in Huntsville
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu*

PROFESSOR YASUHIRO TEZUKA

*Institute of Natural Medicine
Institute of Natural Medicine, University of Toyama,
2630-Sugitani, Toyama 930-0194, Japan
tezuka@innm.u-toyama.ac.jp*

PROFESSOR DAVID E. THURSTON

*Department of Pharmaceutical and Biological Chemistry,
The School of Pharmacy,
University of London, 29-39 Brunswick Square,
London WC1N 1AX, UK
david.thurston@pharmacy.ac.uk*

HONORARY EDITOR**PROFESSOR GERALD BLUNDEN**

*The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com*

ADVISORY BOARD

Prof. Berhanu M. Abegaz
Gaborone, Botswana

Prof. Viqar Uddin Ahmad
Karachi, Pakistan

Prof. Øyvind M. Andersen
Bergen, Norway

Prof. Giovanni Appendino
Novara, Italy

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Lee Banting
Portsmouth, U.K.

Prof. Julie Banerji
Kolkata, India

Prof. Anna R. Bilia
Florence, Italy

Prof. Maurizio Bruno
Palermo, Italy

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Cristina Gracia-Viguera
Murcia, Spain

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. A.A. Leslie Gunatilaka
Tucson, AZ, USA

Prof. Kurt Hostettmann
Lausanne, Switzerland

Prof. Martin A. Iglesias Arteaga
Mexico, D. F, Mexico

Prof. Jerzy Jaroszewski
Copenhagen, Denmark

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Teodoro Kaufman
Rosario, Argentina

Prof. Norbert De Kimpe
Gent, Belgium

Prof. Karsten Krohn
Paderborn, Germany

Prof. Hartmut Laatsch
Gottingen, Germany

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Shoen-Sheng Lee
Taipei, Taiwan

Prof. Francisco Macias
Cadiz, Spain

Prof. Imre Mathe
Szeged, Hungary

Prof. Joseph Michael
Johannesburg, South Africa

Prof. Ermino Murano
Trieste, Italy

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Om Prakash
Manhattan, KS, USA

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Satyajit Sarker
Wolverhampton, UK

Prof. Monique Simmonds
Richmond, UK

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Karen Valant-Vetschera
Vienna, Austria

Prof. Peter G. Waterman
Lismore, Australia

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2010 subscription price: US\$1,695 (Print, ISSN# 1934-578X); US\$1,695 (Web edition, ISSN# 1555-9475); US\$2,095 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Immunomodulatory Action of Monosulfated Triterpene Glycosides from the Sea Cucumber *Cucumaria okhotensis*: Stimulation of Activity of Mouse Peritoneal Macrophages

Dmitry L. Aminin^a, Alexandra S. Silchenko^a, Sergey A. Avilov^a, Vadim G. Stepanov^b and Vladimir I. Kalinin^{a,*}

^aPacific Institute of Bioorganic Chemistry, Vladivostok, 690022, Russian Federation

^bKamchatka Department of the Pacific Institute of Geography, Petropavlovsk-Kamchatsky, 683000, Russian Federation

kalininv@piboc.dvo.ru

Received: April 13th, 2010; Accepted: September 24th, 2010

Six monosulfated triterpene glycosides, frondoside A₁ (**1**), okhotoside B₁ (**2**), okhotoside A₁-1 (**3**), frondoside A (**4**), okhotoside A₂-1 (**5**) and cucumarioside A₂-5 (**6**), isolated from *Cucumaria okhotensis* Levin et Stepanov, stimulate spreading and lysosomal activity of mouse macrophages and ROS-formation in the macrophages. The highest macrophage spreading and stimulation of their lysosomal activity was induced by glycosides **1**, **4** and **6**. All glycosides similarly stimulate ROS formation in macrophages, but glycoside **2** caused minimal stimulation.

Keywords: triterpene glycosides, *Cucumaria okhotensis*, macrophages, ROS formation, spreading, lysosomal activity.

Triterpene glycosides of sea cucumbers possess a wide spectrum of biological activity caused by their membranolytic action [1]. During recent decades their immunomodulatory activity in subtoxic doses has attracted attention. Monosulfated glycosides from the sea cucumbers *Cucumaria japonica* [2–5] and *C. frondosa* [6] stimulate mammalian cell immunity while their activity is effective in doses significantly less than the cytotoxic ones against immune cells [7].

Recently it was found that populations of *C. japonica* in the northern parts of the Sea of Japan and Sea of Okhotsk, including the shallow waters of the Kamchatka Peninsula and North Kuril Islands, really are a series of endemic species that differ from *C. japonica* by several features, including composition of triterpene glycosides [8,9]. Because monosulfated glycosides from *C. japonica* are used for the preparation of immunostimulants [5], it is of interest to investigate the immunostimulatory activity of the glycosides isolated recently from the newly described *C. okhotensis* Levin et Stepanov, found near the south-western shore of Kamchatka Peninsula [10].

The structures of the studied monosulfated glycosides from *C. okhotensis* are presented in Figure 1. Glycosides

1 and **4** were also isolated from *C. frondosa* [11,12], and glycoside **6** from *C. conicospermium* [13]. Because the immunomodulatory action of frondoside A (**4**) has been studied previously [6], we used it as a positive control. To study immunomodulatory activity we chose spreading and lysosomal activity of mouse macrophages, and the formation of reactive oxygen species (ROS) in them.

Macrophage adhesion onto an extracellular matrix, followed by their spreading, is necessary to follow their functional activity. These indexes reflect the initial stage of phagocytosis and macrophage ability for phagocytosis. It is accompanied by changes in macrophage geometric parameters, including cell area and perimeter. The cell may elongate and change shape because of the formation of lamellae and filopodia [14,15].

All the studied glycosides induced an increase of all these indexes on the fourth day after intraperitoneal injection. Macrophages of mice administrated with frondoside A (**4**) had larger geometrical parameters and their shape was more differentiated because of an increase in spreading, elongation, and formation of lamellae and filopodia compared with macrophages of mice from the control group.




Figure 1: Chemical structures of monosulfated triterpene glycosides from *Cucumaria okhotensis*: 1 – frondoside A₁ [9,11]; 2 – okhotoside B₁ [9]; 3 – okhotoside A₁⁻¹ [8]; 4 – frondoside A [9,12]; 5 – okhotoside A₂⁻¹ [8]; 6 – cucumarioside A₂⁻⁵ [9,13].

Table 1: Effect of triterpene glycosides on peritoneal macrophage spreading. The glycosides were administrated by intraperitoneal injection of 0.5 mL at a dose of 0.2 µg/mouse. On the fourth day the mice were killed by perversical dislocation followed by isolation of macrophages that were then stained with the fluorescent probe 5-carboxyfluorescein diacetate and analyzed. The data are presented as $m \pm se$ ($n=100$); *— $p < 0.05$.

Substance	Geometric parameters of macrophages, pixels			
	Cell area	Perimeter	Maximal chord	Shape
Control	92.163 ± 13.111	80.698 ± 9.484	11.050 ± 1.094	0.189 ± 0.016
Fr A ₁ (1)	239.315 ± 25.444*	104.849 ± 7.578*	21.284 ± 1.465*	0.245 ± 0.012*
Okh B ₁ (2)	107.989 ± 15.989	120.333 ± 8.621*	14.774 ± 0.863*	0.075 ± 0.003*
Okh A ₁ ⁻¹ (3)	154.146 ± 18.282*	142.343 ± 12.40*	19.214 ± 1.452*	0.106 ± 0.007*
Fr A (4)	250.125 ± 33.284*	128.659 ± 12.240*	22.472 ± 1.611*	0.210 ± 0.011*
Okh A ₂ ⁻¹ (5)	157.138 ± 22.937*	88.400 ± 8.956	17.493 ± 1.064*	0.285 ± 0.011*
Cucu A ₂ ⁻⁵ (6)	229.608 ± 39.828*	125.411 ± 16.941*	22.592 ± 2.453*	0.206 ± 0.014

Almost all the studied glycosides induced statistically significant increases in cell area, perimeter, maximal chord and integral index of cell shape at an injection dose of 0.2 µg/mouse. Quantitative estimation of the influence of the glycosides on macrophage spreading is presented in Table 1. The most effective glycosides were frondoside A₁ (1), frondoside A (4) and cucumarioside A₂-5 (6). These substances activate spreading 2–2.5 fold greater than the control. Frondoside A was more active than frondoside A₁ and cucumarioside A₂-5.

Lysosomal activity is one of the important markers of physiological and biochemical macrophage status. Almost all the studied glycosides induced a statistically significant increase in lysosomal activity of peritoneal macrophages on the fourth day after a single intraperitoneal injection of 0.2 µg/mouse. The most effective glycosides were frondoside A₁ (1), frondoside A (4) and cucumarioside A₂-5 (6). These compounds activated lysosomal activity approximately 1.5–2.5 times greater than that of control cells (Figure 2). The most active glycoside was frondoside A, whereas the activity of glycoside 6 was similar to that of glycoside 1.




Figure 2: Influence of triterpene glycosides on lysosomal activity of mouse peritoneal macrophages. Glycosides were administrated by intraperitoneal injection of 0.5 mL at a dose 0.2 µg/mouse. On the fourth day the mice were killed by perversical dislocation followed by isolation of macrophages. The lysosomes were stained with acridine orange and analyzed. The data are presented as $m \pm se$ ($n=100$); *— $p < 0.05$.

The ability to generate ROS, singlet oxygen, hydrogen peroxide and the products of their interaction with each other, and NO, the so called “oxidative burst”, is

important for the ability of macrophages to kill ingested microorganisms.

Almost all the studied glycosides induced statistically significant increases of ROS formation in peritoneal macrophages isolated from mice on the fourth day after a single intraperitoneal injection at a dose of 0.2 $\mu\text{g}/\text{mouse}$ (Figure 3). All the glycosides activated lysosomal activity 1.3–1.8 folds greater than control cells. Okhotoside B₁ (**2**) showed minimal activity.

Hence all the studied monosulfated glycosides isolated from *Cucumaria okhotensis* showed immunostimulatory activity of mouse macrophages, as indicated by an increase in macrophage spreading, their lysosomal activity and ROS-formation. This indicates that *C. okhotensis* can be used as a source for immunostimulant preparations.

The activity of the studied glycosides was changed depending on their structure, but not significantly. Glycosides **4** and **5** having a fifth terminal monosaccharide unit (xylose) were more active than their corresponding tetrasaccharide analogs **1** and **2**.




Figure 3: Influence of triterpene glycosides on the ROS-formation in peritoneal mouse macrophages. The glycosides were administrated by intraperitoneal injection of 0.5 mL at a dose of 0.2 $\mu\text{g}/\text{mouse}$. On the fourth day the mice were killed by perversal dislocation followed by isolation of macrophages. The cells were stained with dihydrorhodamine 123 probe and analyzed. The data are presented as $m \pm se$ ($n=100$); *– $p < 0.05$.

Experimental

Triterpene glycosides: The glycosides were isolated from *Cucumaria okhotensis* using common procedures of hydrophobic chromatography of the extracts on teflon powder Polichrom-1, followed by chromatography on a Si gel column and HPLC [8,9]. Structures and purity of individual isolated frondoside A₁ (**1**), okhotoside B₁ (**2**), okhotoside A₁-1 (**3**), frondoside A (**4**), okhotoside A₂-1 (**5**) and cucumarioside A₂-5 (**6**) were confirmed by ¹³C NMR

spectra [8,9]. Individual substances were dissolved in sterile distilled water at a concentration of 1 mg/mL (initial concentration). Forty μL initial glycoside solution were added to 960 μL of distilled water to produce a solution of 40 $\mu\text{g}/\text{mL}$. Ten μL of this solution was added to 990 μL of distilled water to obtain the final concentration of 0.4 $\mu\text{g}/\text{mL}$. The final concentration was used for injection into mice.

Determination of immunomodulatory activity: CBA mice (female, 20 g) were used for *in vivo* testing. The glycosides were intraperitoneally injected (0.5 mL of a glycoside solution in distilled water) at a final glycoside dose of 0.2 $\mu\text{g}/\text{mouse}$. The control mice were injected with distilled water. On the fourth day after injection the mice were killed by perversal dislocation and peritoneal macrophages were isolated by standard procedures. The estimation of immunomodulatory activity was carried out by staining and localization of lysosome in macrophages, determination of reactive oxygen species (ROS) formation in macrophages, and determination of macrophage spreading on an extracellular matrix. Molecular fluorescent probes followed by cell imaging analysis were used. Mice peritoneal liquor (250 μL) containing macrophages was placed onto a microscope cover slip and incubated for 1 h at 37°C. After macrophage adhesion, the cover slip was washed twice with phosphate buffered saline (PBS, pH 7.5). Then 250 μL of a solution of the fluorescent probe was added to the cell monolayer in drops and the cells were incubated at 37°C. To determine macrophage lysosomal activity, a solution of acridine orange (Calbiochem, 100 $\mu\text{g}/\text{mL}$ in PBS) was applied and the cells were incubated for 30 min. To estimate ROS formation, a solution of dihydrorhodamine 123 (Sigma, 100 $\eta\text{g}/\text{mL}$ in PBS containing 0.5 mM of sodium azide) was used. The cells were incubated for 10 min. To estimate macrophage spreading, a solution of 5-carboxyfluorescein diacetate (Molecular Probes, 50 $\mu\text{g}/\text{mL}$ in PBS) was applied. The cells were incubated for 60 min. The cell monolayer was washed twice with PBS. Cover slips were placed on the stage of a fluorescent scanning device composed on the base of an inverted microscope Axiovert 200 (Zeiss, Germany). A luminiscent 75 W Optosource xenon arc lamp and a DAC-controlled monochromator Optoscan (Cairn Research Ltd., UK) were used as light sources for inducing fluorescence at $\lambda_{\text{ex}} = 488 \text{ nm}$; a HQ FITC filter-block (Chroma Technology Corp., USA) and Fluor 40 \times /1.3 oil objective lens (Zeiss, Germany) were used to visualize cell fluorescence. Cell fluorescence images were recorded using a Hamamatsu Orca-ER C4742-95 digital monochrome video camera (Hamamatsu Photonics K.K., Japan) and an IBM-compatible computer having a Fireware data interface.

The intensity of fluorescence of 100 randomly chosen cell images was estimated using an AQM Advance 6 computer program (Kinetic Imaging Ltd., UK) and calculated along with average intensity of fluorescence for each cell, in pixels. Geometric cell parameters (area, perimeter, maximal chord and shape) were also estimated. Each experiment was repeated in triplicate. Average value, standard error, standard deviation and p-values were calculated and plotted on the chart using a SigmaPlot 3.02 computer program (Jandel Scientific, USA).

Acknowledgements – The authors acknowledge the financial support of a Grant of Presidium of the Russian Academy of Science “Basic Researches for Medicine”, the President of the Russian Federation Program for Support of the Leading Scientific Schools Grant No. 3531.2010.4 and the Russian Government Contract No. 02.740.11.0777. The authors very much appreciated the help of Professor John M. Lawrence, University of South Florida (Tampa, FL) for correction of the manuscript.

References

- [1] Kalinin VI, Aminin DL, Avilov SA, Silchenko AS, Stonik VA. (2008) Triterpene glycosides from sea cucumbers (Holothurioidea, Echinodermata), biological activities and functions. In: *Studies in Natural Product Chemistry (Bioactive Natural Products)*, Vol. 35. Atta-ur-Rahman (Ed.). Elsevier Science Publisher, Amsterdam, 135–196.
- [2] Aminin DL, Agafonova IG, Berdyshev EV, Isachenko EG, Avilov SA, Stonik VA. (2001) Immunomodulatory properties of cucumariosides from the edible Far-Eastern holothurian *Cucumaria japonica*. *Journal of Medicinal Food*, 4, 127–135.
- [3] Agafonova IG, Aminin DL, Avilov SA, Stonik VA. (2003) Influence of cucumariosides upon intracellular $[Ca^{2+}]_i$ and lysosomal activity of macrophages. *Journal of Agricultural and Food Chemistry*, 51, 6982–6986.
- [4] Aminin DL, Koy C, Dmitrenok PS, Muller-Hilke B, Koczan D, Arbogast B, Silchenko AS, Kalinin VI, Avilov SA, Stonik VA, Collin PD, Thiersen H-J, Deinzer ML, Glocker MO. (2009) Immunomodulatory effects of holothurian triterpene glycosides on mammalian splenocytes determined by mass spectrometric proteome analysis. *Journal of Proteomics*, 72, 886–906.
- [5] Aminin DL, Pinegin BV, Pichugina LV, Zaporozhets TS, Agafonova IG, Boguslavsky VM, Silchenko AS, Avilov SA, Stonik VA. (2006) Immunomodulatory properties of cumaside. *International Immunopharmacology*, 6, 1070–1082.
- [6] Aminin DL, Agafonova IG, Kalinin VI, Silchenko AS, Avilov SA, Stonik VA, Collin PD, Woodward C. (2008) Immunomodulatory properties of frondoside A, a major triterpene glycoside from the North Atlantic commercially harvested sea cucumber *Cucumaria frondosa*. *Journal of Medicinal Food*, 11, 443–453.
- [7] Aminin DL, Silchenko AS, Avilov SA, Stepanov VG, Kalinin VI. (2009) Cytotoxic action of triterpene glycosides from sea cucumbers from the genus *Cucumaria* on mouse spleen lymphocytes. Inhibition of nonspecific esterase. *Natural Product Communications*, 4, 773–776.
- [8] Silchenko AS, Avilov SA, Kalinin VI, Stonik VA, Kalinovskiy AI, Dmitrenok PS, Stepanov VG. (2007) Monosulfated triterpene glycosides from *Cucumaria okhotensis* Levin et Stepanov – new sea cucumber species from the Sea of Okhotsk. *Bioorganicheskaya Khimiya*, 33, 81–90.
- [9] Silchenko AS, Avilov SA, Kalinin VI, Kalinovskiy AI, Dmitrenok PS, Fedorov SN, Stepanov VG, Dong Z, Stonik VA. (2008) Constituents of the sea cucumber *Cucumaria okhotensis*. Structures of okhotosides B₁–B₃ and cytotoxic activities of some glycosides from this species. *Journal of Natural Products*, 71, 351–356.
- [10] Levin VS. (2003) *Cucumaria okhotensis* – a new species of holothurians (Echinodermata: Holothuriodea) from the Sea of Okhotsk. *Biologiya Morya*, 29, 202–205.
- [11] Avilov SA, Kalinin VI, Drozdova OA, Kalinovskiy AI, Stonik VA, Gudimova EN. (1993) Triterpene glycosides from the sea cucumber *Cucumaria frondosa*. *Khimiya Prirodnih Soedinenii*, 2, 49–52.
- [12] Girard M, Belanger J, ApSimon JW, Garneau F-X, Harvey C, Brisson J-R. Frondoside A. (1990) A novel triterpene glycoside from the holothurian *Cucumaria frondosa*. *Canadian Journal of Chemistry*, 68, 11–18.
- [13] Avilov SA, Antonov AS, Silchenko AS, Kalinin VI, Kalinovskiy AI, Dmitrenok PS, Stonik VA, Riguera R, Jimenes C. (2003) Triterpene glycosides from the Far Eastern sea cucumber *Cucumaria conicospermum*. *Journal of Natural Products*, 66, 910–916.
- [14] Mosser DM, Edwards JP. (2008) Exploring the full spectrum of macrophage activation. *Nature Reviews in Immunology*, 8, 958–969.
- [15] Martinez FO, Sica A, Mantovani A, Locati M. (2008) Macrophage activation and polarization. *Frontiers in Bioscience*, 13, 453–461.

Inhibition of Protein Tyrosine Phosphatase 1β by Hispidin Derivatives Isolated from the Fruiting Body of <i>Phellinus linteus</i>	1927
Yeon Sil Lee, Il-Jun Kang, Moo Ho Won, Jae-Yong Lee, Jin Kyu Kim and Soon Sung Lim	
A New Azafluorenone from the Roots of <i>Polyalthia cerasoides</i> and its Biological Activity	1931
Kanchana Pumsalid, Haruthai Thaisuchat, Chatchanok Loetchutinat, Narong Nuntasaeen, Puttinan Meepowpan and Wilart Pompimon	
Evaluation of Antiviral Activities of Curcumin Derivatives against HSV-1 in Vero Cell Line	1935
Keivan Zandi, Elissa Ramedani, Khosro Mohammadi, Saeed Tajbakhsh, Iman Deilami, Zahra Rastian, Moradali Fouladvand, Forough Yousefi and Fatemeh Farshadpour	
Hyloglyceride and Hylodiglyceride: Two New Glyceride Derivatives from <i>Hylodendron gabunensis</i>	1939
Awazi Tengu Nyongha, Hidayat Hussain, Etienne Dongo, Ishtiaq Ahmed and Karsten Krohn	
Chemical Composition and Bioactivities of the Marine Alga <i>Isochrysis galbana</i> from Taiwan	1941
Chi-Cheng Yu, Hsiao-Wei Chen, Mao-Jing Chen, Yu-Ching Chang, Shih-Chang Chien, Yueh-Hsiung Kuo, Feng-Ling Yang, Shih-Hsiung Wu, Jie Chen, Hsiao-Hui Yu and Louis Kuop-Ping Chao	
An Efficient Protocol for High-frequency Direct Multiple Shoot Regeneration from Internodes of Peppermint (<i>Mentha x piperita</i>)	1945
Sanjog T. Thul and Arun K. Kukreja	
Essential Oil Yield and Chemical Composition Changes During Leaf Ontogeny of Palmarosa (<i>Cymbopogon martinii</i> var. <i>motia</i>)	1947
Bhaskaruni R. Rajeswara Rao, Dharmendra K. Rajput, Rajendra P. Patel and Somasi Purnanand	
Essential Oil Composition of Four Endemic <i>Ferulago</i> Species Growing in Turkey	1951
Ceyda Sibel Kılıç, Ayşe Mine Gençler Özkan, Betül Demirci, Maksut Coşkun and Kemal Hüsnü Can Başer	
Essential Oils of <i>Daucus carota</i> subsp. <i>carota</i> of Tunisia Obtained by Supercritical Carbon Dioxide Extraction	1955
Hanen Marzouki, Abdelhamid Khaldi, Danilo Falconieri, Alessandra Piras, Bruno Marongiu, Paola Moliccotti and Stefania Zanetti	
Oil Constituents of <i>Artemisia nilagirica</i> var. <i>septrionalis</i> Growing at Different Altitudes	1959
Flora Haider, Narendra Kumar, Ali Arif Naqvi and Guru Das Bagchi	
Volatile Oil Composition of <i>Pogostemon heyneanus</i> and Comparison of its Composition with Patchouli Oil	1961
Ramar Murugan, Gopal Rao Mallavarapu, Kyathsandra Venkataramaiah Padmashree, Ramachandra Raghavendra Rao and Christus Livingstone	
Chemical Composition of Volatile Oils of <i>Aquilaria malaccensis</i> (Thymelaeaceae) from Malaysia	1965
Saiful Nizam Tajuddin and Mashitah M. Yusoff	
Chemical Composition and Phytotoxic Effects of Essential Oils from Four <i>Teucrium</i> Species	1969
Laura De Martino, Carmen Formisano, Emilia Mancini, Vincenzo De Feo, Franco Piozzi, Daniela Rigano and Felice Senatore	
Chemical Constituents and Larvicidal Activity of <i>Hymenaea courbaril</i> Fruit Peel	1977
José Cláudio D. Aguiar, Gilvandete M. P. Santiago, Patrícia L. Lavor, Helenicy N. H. Veras, Yana S. Ferreira, Michele A. A. Lima, Ângela M. C. Arriaga, Telma L. G. Lemos, Jefferson Q. Lima, Hugo C. R. de Jesus, Pérciles B. Alves and Raimundo Braz-Filho	
Caryophyllene Oxide-rich Essential Oils of Lithuanian <i>Artemisia campestris</i> ssp. <i>campestris</i> and Their Toxicity	1981
Asta Judzentiene, Jurga Budiene, Rita Butkiene, Eugenija Kupcinskiene, Isabelle Laffont-Schwob and Véronique Masotti	
Comparison of Antibacterial Activity of Natural and Hydroformylated Essential Oil of <i>Thymus capitatus</i> Growing Wild in North Sardinia with Commercial <i>Thymus</i> Essential Oils	1985
Marianna Usai, Marzia Foddai, Barbara Sechi, Claudia Juliano and Mauro Marchetti	
Composition and Chemical Variability of the Leaf Oil from Corsican <i>Juniperus thurifera</i> Integrated Analysis by GC(RI), GC-MS and ¹³C NMR	1991
Josephine Ottavioli, Joseph Casanova and Ange Bighelli	
Combined Analysis by GC (RI), GC-MS and ¹³C NMR of the Supercritical Fluid Extract of <i>Abies alba</i> Twigs	1995
Emilie Duquesnoy, Bruno Marongiu, Vincent Castola, Alessandra Piras, Silvia Porcedda and Joseph Casanova	
<u>Review/Account</u>	
Eugenol: A Natural Compound with Versatile Pharmacological Actions	1999
Kannissery Pramod, Shahid H. Ansari and Javed Ali	

Natural Product Communications

2010

Volume 5, Number 12

Contents

<u>Original Paper</u>	<u>Page</u>
Anticonvulsant Activity of the Linalool Enantiomers and Racemate: Investigation of Chiral Influence Damião P. de Sousa, Franklin F. F. Nóbrega, Camila C. M. P. Santos and Reinaldo N. de Almeida	1847
Kinetic Analysis of Genipin Degradation in Aqueous Solution Paul Slusarewicz, Keng Zhu and Tom Hedman	1853
Microbial Transformation of Marine Halogenated Sesquiterpenes Aurelio San Martin, Juana Rovirosa, Alvaro Carrasco, Silvia Orejarena, Jorge Soto-Delgado, Renato Contreras and M. Cristina Chamy	1859
Two New Guaianolides from <i>Amberboa ramosa</i> Muhammad Ibrahim, Rehan Khan and Abdul Malik	1865
Antiplasmodial and Cytotoxic Activities of Drimane Sesquiterpenes from <i>Canella winterana</i> Mary H. Grace, Carmen Lategan, Flaubert Mbeunkui, Rocky Graziose, Peter J. Smith, Ilya Raskin and Mary Ann Lila	1869
Three New 18-Oxygenated <i>ent</i>-Kaurane Diterpenoids from <i>Isodon leucophyllus</i> Hai Bo Zhang, Jian Xin Pu, Yong Zhao, Fei He, Wei Zhao, Li Guang Lou, Wei Lie Xiao and Han Dong Sun	1873
Immunomodulatory Action of Monosulfated Triterpene Glycosides from the Sea Cucumber <i>Cucumaria okhotensis</i>: Stimulation of Activity of Mouse Peritoneal Macrophages Dmitry L. Aminin, Alexandra S. Silchenko, Sergey A. Avilov, Vadim G. Stepanov and Vladimir I. Kalinin	1877
Three New Aaptamines from the Marine Sponge <i>Aaptos</i> sp. and Their Proapoptotic Properties Larisa K. Shubina, Tatyana N. Makarieva, Sergey A. Dyshlovoy, Sergey N. Fedorov, Pavel S. Dmitrenok and Valentin A. Stonik	1881
Isolation and Characterization of Crotosparsamide, a New Cyclic Nonapeptide from <i>Croton sparsiflorus</i> Rashad Mehmood and Abdul Malik	1885
Two New Lavandulyl Flavonoids from <i>Sophora flavescens</i> Dan Liu, Xiulan Xin, Dong-hai Su, Junying Liu, Qing Wei, Bo Li and Jian Cui	1889
Biotransformation of Naringenin to Eriodictyol by <i>Saccharomyces cerevisiae</i> Functionally Expressing Flavonoid 3' Hydroxylase Ilef Limem-Ben Amor, Alain Hehn, Emmanuel Guedon, Kamel Ghedira, Jean-Marc Engasser, Leila Chekir-Ghedira and Mohamed Ghoul	1893
Two New 3-C-Carboxylated Flavones from the Rhizomes of <i>Caragana conferta</i> Rehan Khan, Abdul Malik, Shazia Yasmeen and Nighat Afza	1899
Kaempferol Glycosides in the Flowers of Carnation and their Contribution to the Creamy White Flower Color Tsukasa Iwashina, Masa-atsu Yamaguchi, Masayoshi Nakayama, Takashi Onozaki, Hiroyuki Yoshida, Shuji Kawanobu, Hiroshi Ono and Masachika Okamura	1903
Factors Influencing Glabridin Stability Mingzhang Ao, Yue Shi, Yongming Cui, Wentao Guo, Jing Wang and Longjiang Yu	1907
Effect of Different Strains of <i>Agrobacterium rhizogenes</i> and Nature of Explants on <i>Plumbago indica</i> Hairy Root Culture with Special Emphasis on Root Biomass and Plumbagin Production Moumita Gangopadhyay, Saikat Dewanjee, Somnath Bhattacharyya and Sabita Bhattacharya	1913
Fujianmycin C, A Bioactive Angucyclinone from a Marine Derived <i>Streptomyces</i> sp. B6219 Muna Ali Abdalla, Elisabeth Helmke and Hartmut Laatsch	1917
Dioscorealide B from the Traditional Thai Medicine Hua-Khao-Yen Induces Apoptosis in MCF-7 Human Breast Cancer Cells via Modulation of Bax, Bak and Bcl-2 Protein Expression Jiraporn Saekoo, Potchanapond Graidist, Wilairat Leeanansaksiri, Chavaboon Dechsukum and Arunporn Itharat	1921

Continued inside backcover